

On some integral classes of integral operators¹

Virgil Pescar

Abstract

Let A be the class of the functions f which are analytic in the unit disk $U = \{z \in C; |z| < 1\}$ and $f(0) = f'(0) - 1 = 0$. The object of the present paper is to derive univalence conditions of certain integral operators for $f(z) \in A$ and $f(z)$ has the form: $f(z) = z + \sum_{k=3}^{\infty} a_k z^k$.

2000 Mathematics Subject Classification: 30C45.

Key words: Univalent, integral operator.

1 Introduction

Let A be the class of the functions $f(z)$ which are analytic in the unit disk $U = \{z \in C : |z| < 1\}$ and $f(0) = f'(0) - 1 = 0$.

We denote by S the class of the functions $f(z) \in A$ which are univalent in U .

In this paper we consider the integral operators

$$(1.1) \quad F_{\alpha}(z) = \int_0^z [f'(u)]^{\alpha} du$$

¹Received 29 August 2007

Accepted for publication (in revised form) 4 January 2008

$$(1.2) \quad H_{\beta,\gamma}(z) = \left\{ \beta \int_0^z u^{\beta-1} [f'(u)]^\gamma du \right\}^{\frac{1}{\beta}}$$

$$(1.3) \quad L_\beta(z) = \left[\beta \int_0^z u^{\beta-1} [f'(u)] du \right]^{\frac{1}{\beta}}$$

2 Preliminary Results

We need the following theorems.

Lemma 2.1. [1]. *If $f(z) \in A$ satisfies*

$$(2.1) \quad (1 - |z|^2) \left| \frac{z f''(z)}{f'(z)} \right| \leq 1, \quad z \in U$$

then $f(z) \in S$.

Theorem 2.2. [3]. *Let α be a complex number, $\operatorname{Re} \alpha > 0$ and $f(z) \in A$. If*

$$(2.2) \quad \frac{1 - |z|^{2\operatorname{Re} \alpha}}{\operatorname{Re} \alpha} \left| \frac{z f''(z)}{f'(z)} \right| \leq 1,$$

for all $z \in U$, then for any complex number β , $\operatorname{Re} \beta \geq \operatorname{Re} \alpha$ the function

$$(2.3) \quad F_\beta(z) = \left[\beta \int_0^z u^{\beta-1} f'(u) du \right]^{\frac{1}{\beta}}$$

is in the class S .

Theorem 2.3. [2]. *If the function $g(z)$ is regular in U and $|g(z)| < 1$ in U , then for all $\xi \in U$ and $z \in U$ the following inequalities hold:*

$$(2.4) \quad \left| \frac{g(\xi) - g(z)}{1 - \overline{g(z)}g(\xi)} \right| \leq \left| \frac{\xi - z}{1 - \overline{z}\xi} \right|,$$

$$(2.5) \quad |g'(z)| \leq \frac{1 - |g(z)|^2}{1 - |z|^2},$$

the equalities hold only in the case $g(z) = \frac{\epsilon(z+u)}{1+\bar{u}z}$, where $|\epsilon| = 1$ and $|u| < 1$.

Remark 2.4. [2] For $z = 0$, from inequality (2.4). We have

$$(2.6) \quad \left| \frac{g(\xi) - g(0)}{1 - \overline{g(0)}g(\xi)} \right| \leq |\xi|$$

and, hence

$$(2.7) \quad |g(\xi)| \leq \frac{|\xi| + |g(0)|}{1 + |g(0)||\xi|}.$$

Considering $g(0) = a$ and $\xi = z$,

$$(2.8) \quad |g(z)| \leq \frac{|z| + |a|}{1 + |a||z|}$$

for all $z \in U$.

3 Main Results

Theorem 3.1. Let α be a complex number and $f(z) \in A$,

$$f(z) = z + \sum_{k=3}^{\infty} a_k z^k. \text{ If}$$

$$(3.1) \quad \left| \frac{f''(z)}{f'(z)} \right| < 1, \quad z \in U$$

and

$$(3.2) \quad |\alpha| \leq 4$$

then the function

$$(3.3) \quad F_\alpha(z) = \int_0^z [f'(u)]^\alpha du$$

is in the class S .

Proof. The function $F_\alpha(z)$ is regular in U . Let us consider the function

$$(3.4) \quad p(z) = \frac{1}{|\alpha|} \frac{F_\alpha''(z)}{F_\alpha'(z)}$$

where the constant $|\alpha|$ satisfies the inequality (3.2).

The function $p(z)$ is regular in U . From (3.4) and (3.3) we obtain

$$(3.5) \quad p(z) = \frac{\alpha}{|\alpha|} \frac{f''(z)}{f'(z)}.$$

Using (3.1) and (3.5) we get

$$(3.6) \quad |p(z)| \leq 1, \quad z \in U$$

and we have $p(0) = 0$.

By Remark 2.4 we have

$$(3.7) \quad |p(z)| \leq |z|, \quad z \in U.$$

From (3.4) and (3.7) we obtain

$$(3.8) \quad \frac{1}{|\alpha|} \left| \frac{F''_{\alpha}(z)}{F'_{\alpha}(z)} \right| \leq |z|, \quad z \in U$$

and

$$(3.9) \quad (1 - |z|^2) \left| \frac{zF''_{\alpha}(z)}{F'_{\alpha}(z)} \right| \leq |\alpha| \max_{|z|<1} (1 - |z|^2) |z|^2.$$

Because $\max_{|z|<1} (1 - |z|^2) |z|^2 = \frac{1}{4}$, from (3.9) and (3.2) we get

$$(3.10) \quad (1 - |z|^2) \left| \frac{zF''_{\alpha}(z)}{F'_{\alpha}(z)} \right| \leq 1, \quad z \in U.$$

By Lemma 2.1 it results that the function $F_{\alpha}(z) \in S$.

Theorem 3.2. *Let γ be a complex number and the function $f(z) \in A$,*

$f(z) = z + \sum_{k=3}^{\infty} a_k z^k$. If

$$(3.11) \quad \left| \frac{f''(z)}{f'(z)} \right| < 1, \quad z \in U$$

and

$$(3.12) \quad |\gamma| \leq 4$$

then for any complex number β , $\operatorname{Re} \beta \geq 1$ the function

$$(3.13) \quad H_{\beta,\gamma}(z) = \left\{ \beta \int_0^z u^{\beta-1} [f'(u)]^\gamma du \right\}^{\frac{1}{\beta}}$$

is in the class S .

Proof. Let us consider the function

$$(3.14) \quad g(z) = \int_0^z [f'(u)]^\gamma du.$$

The function

$$(3.15) \quad p(z) = \frac{1}{|\gamma|} \frac{g''(z)}{g'(z)},$$

where the constant $|\gamma|$ satisfies the inequality (3.12), is regular in U .

From (3.15) and (3.14) we obtain

$$(3.16) \quad p(z) = \frac{\gamma}{|\gamma|} \frac{f''(z)}{f'(z)}.$$

and using (3.11) we have

$$|p(z)| \leq 1, \quad z \in U$$

Remark 2.4 applied to the function $p(z)$ give

$$(3.17) \quad \frac{1}{|\gamma|} \left| \frac{g''(z)}{g'(z)} \right| \leq |z|, \quad z \in U$$

and, hence

$$(3.18) \quad (1 - |z|^2) \left| \frac{zg''(z)}{g'(z)} \right| \leq |\gamma| \max_{|z|<1} (1 - |z|^2) |z|^2.$$

From (3.18) and (3.12) we obtain

$$(3.19) \quad (1 - |z|^2) \left| \frac{zg''(z)}{g'(z)} \right| \leq 1, \quad z \in U.$$

By Theorem 2.2 for $Re \alpha = 1$, it results that $H_{\beta, \gamma}(z) \in S$.

Theorem 3.3. *Let β a complex number, $Re \beta \geq 1$ and $f(z) \in A$, $f(z) = z + a_3z^3 + \dots$, $\frac{f(z)}{z} \neq 0$, $z \in U$. If*

$$(3.20) \quad \left| \frac{f''(z)}{f'(z)} \right| \leq 4, \quad z \in U$$

then the function

$$(3.21) \quad L_\beta(z) = \left[\beta \int_0^z u^{\beta-1} [f'(u)] du \right]^{\frac{1}{\beta}}$$

is in the class S .

Proof. Let us consider the function

$$g(z) = \frac{1}{4} \frac{f''(z)}{f'(z)}$$

which is regular in U . Remark 2.4 applied to the function $g(z)$ give

$$(3.22) \quad \frac{1}{4} \left| \frac{f''(z)}{f'(z)} \right| \leq |z|, \quad z \in U$$

and, hence, we obtain

$$(3.23) \quad (1 - |z|^2) \left| \frac{zf''(z)}{f'(z)} \right| \leq 4 \max_{|z|<1} (1 - |z|^2) |z|^2, \quad z \in U$$

Since $\max_{|z|<1} (1 - |z|^2) |z|^2 = \frac{1}{4}$, from (3.23) we have

$$(3.24) \quad (1 - |z|^2) \left| \frac{zf''(z)}{f'(z)} \right| \leq 1, \quad z \in U.$$

From (3.24) and Theorem 2.2 for $Re \alpha = 1$, we obtain $F_\beta(z) \in S$.

References

- [1] J. Becker, *Löwnersche Differentialgleichung und quasikonform fortsetzbare schlichte Functionen*, J. Reine Angew. Math., 255 (1972), 23-43.
- [2] Z. Nehari, *Conformal mapping*, Mc Graw-Hill Book Co., Inc., New York, Toronto, London, 1952 .
- [3] N. N. Pascu, *An improvement of Beckers univalence criterion*, Proceedings of the Commemorative Session Simion Stoilow (Braşov), 1987, University of Braşov, pp. 43-48.
- [4] V. Pescar, *New univalence criteria*, Transilvania University of Braşov, 2002.
- [5] C. Pommerenke, *Univalent functions*, Vanderhoeck Ruprecht in Göttingen, 1975.

”Transilvania” University of Braşov
Faculty of Mathematics and Computer Science
Department of Mathematics
2200 Braşov, Romania
E-mail: virgilpescar@unitbv.ro